

California Bioscience

83103 Avenue 48, Ste.1B #204 Coachella, CA 92236 USA Phone : +1.6268339877 Email : info@cali-bio.com

Product Datasheet

Product Name	Bone Morphogenetic Protein-7 Human Recombinant, CHO
Cata No	CB500178
Source	Chinese Hamster Ovarian Cells
Synonyms	Osteogenic Protein 1, BMP-7.

Description

The bone morphogenetic proteins (BMPs) are a family of secreted signaling molecules that can induce ectopic bone growth. Many BMPs are part of the transforming growth factor-beta (TGFB) superfamily. BMPs were originally identified by an ability of demineralized bone extract to induce endochondral osteogenesis in vivo in an extraskeletal site. Based on its expression early in embryogenesis, the BMP encoded by this gene has a proposed role in early development. In addition, the fact that this BMP is closely related to BMP5 and BMP7 has lead to speculation of possible bone inductive activity.

N-TERMINAL---Human BMP-2 (Met 1 – Arg 282) Human BMP-7 (Ser 293 – Arg 431)---C-TERMINAL. The DNA sequence encoding the human BMP-2 signal peptide and propeptide (1~282 amino acid) fused to the human rhBMP-7 mature chain (293~431 amino acid) was expressed in a Chinese hamster ovary cell line. The mature recombinant BMP-7 generated by the proteolytic removal of the signal peptide and propetide contains 139 amino acid residues. The glycosylation of BMP-7 increases the molecular mass and the glycosylated proteins migrate as 25 ~ 40 kDa in SDS-PAGE under non-reducing conditions.

BMP-7 is purified by proprietary chromatographic techniques.

Physical Appearance

Sterile Filtered White lyophilized (freeze-dried) powder.

Biological Activity

Measured in alkaline phosphatase activity assay using MC3T3-E1 cells. The ED_{50} for this effect is < 100 ng/mL.

Purity

Greater than 97.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.

Formulation

BMP-7 was lyophilized from a concentrated (1mg/ml) sterile solution containing 1% sucrose, 1.2% mannitol, 20mM glycine, and 0.005% tween 20 pH-4.

Stability

Lyophilized BMP-7 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution BMP 7 Human should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).

Please prevent freeze-thaw cycles.